

Recombinant Human Acidic Fibroblast Growth Factor, 2-155a.a.

#R00105

Size: 10 µg/100 µg/500 µg

Stability & Storage

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.



Warning: This product is intended for scientific research only. It is not to be used for medical or diagnostic procedures in humans or animals, nor as a food, cosmetic, or household product.

AA Sequence

AEGEITTF TALTEKFNLP PGNYKKPKLLYCSNGGHFLR
ILPDGTV DGTDRSDQH IQLQLSAESVGEVYIKSTET
GQYLAMD TDGLLYGSQTPNEECLFLERLEENHYNTYI
SKKHAEKNWFVGLKKN GSCKRGP RTHYGQKAILFLP
LPVSSD

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Product Specification

Accession	P05230
GeneID	2246
Species	Human
Source	Escherichia coli.
Molecular Weight	Approximately 17.3 kDa, a single non-glycosylated polypeptide chain containing 154 amino acids.
Purity	> 97 % by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active when compared to standard. The ED ₅₀ as determined by a cell proliferation assay using murine balb/c 3T3 cells is less than 0.5 ng/ml, corresponding to a specific activity of > 2.0 × 10 ⁶ IU/mg in the presence of 10

Physical Appearance Formulation	µg/ml of heparin. Sterile Filtered White lyophilized (freeze-dried) powder. Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4, with 2 mM EDTA, 0.5 mM DTT, 5 % Trehalose.
Endotoxin	Less than 0.1 EU/µg of rHuaFGF, 2-155a.a. as determined by LAL method.