

Recombinant Murine Interleukin-1 beta

#R00215

Size: 10 µg/100 µg/500 µg

Stability & Storage

- Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
- 12 months from date of receipt, -20 to -70 °C as supplied.
 - 1 month, 2 to 8 °C under sterile conditions after reconstitution.
 - 3 months, -20 to -70 °C under sterile conditions after reconstitution.



Warning: This product is intended for scientific research only. It is not to be used for medical or diagnostic procedures in humans or animals, nor as a food, cosmetic, or household product.

Background

Interleukin-1beta (IL-1β) is a non-secreted proinflammatory cytokine produced mainly by activated macrophages, as well as neutrophils, epithelial cells, and endothelial cells. It possesses metabolic, physiological, haematopoietic activities, and plays one of the central roles in the regulation of the immune responses. Both IL-1α and IL-1β binds to the same receptor and has similar but not identical biological properties. The mature mouse IL1β shares 90 % amino acid sequence identity with cotton rat and rat and 65 %-78 % identity with canine, human and rhesus IL1β.

AA Sequence

VPIRQLHYRLRDEQQKSLVLSDPYELKALHLNGQNIN
QQVIFSMSFVQGEPsNDKIPVALGLKGKNLYLSCVMK
DGTPTLQLESVDPKQYPKKKMEKRFVFNKIEVKSKVE
FESAEPNWWYISTSQAEHKPVFLGNNSGQDIIDFTME

SVSS

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20°C. Further dilutions should be made in appropriate buffered solutions.

Product Specification

Synonyms	Catabolin, Leukocyte Endogenous Mediator, LEM, Lymphocyte-activating factor , LAF, Mononuclear Cell Factor, MCF , Endogenous Pyrogen, EP
Accession	P10749
GeneID	16176
Species	Murine
Source	Escherichia coli.
Molecular Weight	Approximately 17.5 kDa, a single non-glycosylated polypeptide chain containing 152 amino acids.
Purity	> 96 % by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active when compared to standard. The ED ₅₀ as determined by a cell

	proliferation assay using murine D10S cells is less than 2 pg/ml, corresponding to a specific activity of $> 5.0 \times 10^8$ IU/mg.
Physical Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS, pH 7.4, with 5 % trehalose, 0.02 % Tween-20.
Endotoxin	Less than 1 EU/ μ g of rMull-1 β as determined by LAL method.