

Recombinant Murine TNF-related apoptosis-inducing Ligand/TNFSF10

#R00247

Size: 10 µg/100 µg/500 µg

Stability & Storage

- Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
- 12 months from date of receipt, -20 to -70 °C as supplied.
 - 1 month, 2 to 8 °C under sterile conditions after reconstitution.
 - 3 months, -20 to -70 °C under sterile conditions after reconstitution.

**Warning:** This product is intended for scientific research only. It is not to be used for medical or diagnostic procedures in humans or animals, nor as a food, cosmetic, or household product.

AA Sequence

MPRGGRPQKVAAHITGITRRSNSALIPISKDGKTLGQ
KIESWESSRKGHSFLNHVLFRNGELVIEQEGLYIYS
QTYFRFQEAEDASKMVSCKDKVRTKQLVQYIYKYTSY
PDPIVLMKSARNSCWSRDAEYGLYSIQGGGLFELKKK
DRIFVSVTNEHLMDLQDEASFFGAFLIN

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Product Specification

Accession	P50592
GeneID	22035
Species	Murine
Source	Escherichia coli.
Molecular Weight	Approximately 20.2 kDa, a single non-glycosylated polypeptide chain containing 175 amino acids.
Purity	> 95 % by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active when compared to standard. The ED ₅₀ as determined by a cytotoxicity assay using murine L929 cells is less than 0.5 ng/ml, corresponding to a specific activity of > 2.0 × 10 ⁶ IU/mg in the presence of actinomycin D.

Physical Appearance	Sterile Filtered White Lyophilized (freeze-dried) powder.
Formulation	Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4, with 3 mM DTT.
Endotoxin	Less than 0.1 EU/µg of rMuTRAIL/TNFSF10 as determined by LAL method.