

Recombinant Murine Nephroblastoma-overexpressed Gene Protein Homolog

#R00263

Size: 5 µg/100 µg/500 µg

Stability & Storage

- Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
- 12 months from date of receipt, -20 to -70 °C as supplied.
 - 1 month, 2 to 8 °C under sterile conditions after reconstitution.
 - 3 months, -20 to -70 °C under sterile conditions after reconstitution.



Warning: This product is intended for scientific research only. It is not to be used for medical or diagnostic procedures in humans or animals, nor as a food, cosmetic, or household product.

AA Sequence

QVSASLRCPSPKPCPSISPTCAPGVRSVLDGCSC
CPVCARQRGESCSEMRPCDQSSGLYCDRSADPNN
QTGICMVPEGDNCVFDGVIYRNGEKFEPCNCQYFCTC
RDGQIGCLPRCQLDVLLPGPDCPAPRKVAVPGECCE
KWTCGSDEQGTQGTLGGLALPAYRPEATVGVEVSD
SSINCIEQTTEWSACSKSCGMGVSTRVTNRNRQCEM
VKQTRLCIVRPCEQEPEEVTDKKGKKCLRTKKSLKAI
HLQFENCTSLYTYKPRFCGVCS DGRCCTPHNTKTIQ
VEFQCLPGEIHKPVMVIGTCTCYSNCPQNNEAFLQD
LELKTSRGEI

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further

dilutions should be made in appropriate buffered solutions.

Product Specification

Accession	Q64299
GeneID	18133
Species	Murine
Source	Escherichia coli.
Molecular Weight	Approximately 36.4 kDa, a single non-glycosylated polypeptide chain containing 333 amino acids.
Purity	> 95 % by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active when compared to standard. The ED ₅₀ as determined by a cell proliferation assay using murine Balb/c 3T3 cells is less than 1.0 µg/ml, corresponding to a specific activity of > 1000 IU/mg.
Physical	Sterile Filtered White Lyophilized

Appearance	(freeze-dried) powder.
Formulation	Lyophilized from a 0.2 μm filtered concentrated solution in 20 mM Tris-HCl, pH 8.6, 150 mM NaCl.
Endotoxin	Less than 0.1 EU/μg of rMuNOV as determined by LAL method.