

Recombinant Murine Growth Differentiation Factor 7/Bone Morphogenetic Protein-12

#R00284

Size: 10 µg/100 µg/500 µg

Stability & Storage

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.



Warning: This product is intended for scientific research only. It is not to be used for medical or diagnostic procedures in humans or animals, nor as a food, cosmetic, or household product.

Background

Growth/differentiation factors (GDF-1 to GDF-15) are members of the BMP family of TGF-beta superfamily proteins. They are produced as inactive preproteins which are then cleaved and assembled into active secreted homodimers. GDF dimers are disulfide-linked with the exception of GDF-3 and -9. GDF proteins are important during embryonic development, particularly in the skeletal, nervous, and muscular systems.

AA Sequence

TALAGTRGAQGSGGGGGGGGGGGGGGGGGGGGA
GRGHGRRGRSRCRSLHVDKELGWDDWIIAPLD
YEAYHCEGVCDPLRSHLEPTNHAIQTLLNSMAPDA
APASCCVPARLSPISILYIDAANNVVKQYEDMVVEAC
GCR

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom.

Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Product Specification

Accession	P43029
GeneID	238057
Species	Murine
Source	Escherichia coli.
Molecular Weight	Approximately 29.8 kDa, a homodimeric protein consisting of two 146 amino acid non-glycosylated polypeptide chains.
Purity	> 95 % by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active when compared to standard. The ED ₅₀ as determined by inducing alkaline phosphatase production of murine ATDC5 cells is less than 0.5 µg/ml, corresponding to a specific activity of > 2000

	IU/mg.
Physical	Sterile Filtered White lyophilized
Appearance	(freeze-dried) powder.
Formulation	Lyophilized from a 0.2 µm filtered concentrated solution in 30 % Acetonitrile and 0.1 % TFA.
Endotoxin	Less than 0.1 EU/µg of rMuGDF-7/BMP-12 as determined by LAL method.