

Recombinant Canine Interferon-gamma

#R00295

Size: 100 µg/500 µg

Stability & Storage

- Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
- 12 months from date of receipt, -20 to -70 °C as supplied.
 - 1 month, 2 to 8 °C under sterile conditions after reconstitution.
 - 3 months, -20 to -70 °C under sterile conditions after reconstitution.



Warning: This product is intended for scientific research only. It is not to be used for medical or diagnostic procedures in humans or animals, nor as a food, cosmetic, or household product.

AA Sequence

QAMFFKEIENLKEYFNASNPDVSDGGSLFVDILKKWR
EESDKTIIQSQIVSFYKLFDNFKDNQIIQRSMDTIKED
MLGKFLNSSTSKREDFLKLIQIPVNDLQVQRKAINELI
KVMNDLSPRSNLRKRKRSQLFRGRRASK

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Product Specification

Accession	P42161
GeneID	403801
Species	Canine
Source	Escherichia coli.
Molecular Weight	Approximately 16.9 kDa, a single non-glycosylated polypeptide chain containing 143 amino acids.
Purity	> 97 % by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active when compared to standard. The ED ₅₀ as determined by an anti-viral assay using A-72 canine fibroma cells infected with vesicular stomatitis virus (VSV) is less than 2.0 ng/ml, corresponding to a specific activity of > 5.0 × 10 ⁵

Physical Appearance Formulation	IU/mg. Sterile Filtered White lyophilized (freeze-dried) powder. Lyophilized from a 0.2 µm filtered concentrated solution in 25 mM Sodium Succinate, pH 5.0, 60 mM NaCl, with 0.1 % Tween-80.
Endotoxin	Less than 0.1 EU/µg of rCalFN-γ as determined by LAL method.