

Recombinant Rat Interleukin-5

#R00303

Size: 10 µg/100 µg/500 µg

Stability & Storage

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.

- 1 month, 2 to 8 °C under sterile conditions after reconstitution.

- 3 months, -20 to -70 °C under sterile conditions after reconstitution.



Warning: This product is intended for scientific research only. It is not to be used for medical or diagnostic procedures in humans or animals, nor as a food, cosmetic, or household product.

Background

IL-5, also named B-cell differentiation factor I, eosinophil differentiation factor and TRF, is belonging to the cytokine family and the IL-5 gene is in close proximity to the genes encoding IL-3, IL-4, and granulocyte-macrophage colony-stimulating factor (GM-CSF), which are often co-expressed in TH2 cells. Through binding to the IL-5 receptor, IL-5 stimulates B cell growth and increases immunoglobulin secretion. It is also a key mediator in eosinophil activation. Interleukin-5 has long been associated with the cause of several allergic diseases including allergic rhinitis and asthma. Rat IL-5 is a 132-amino acid (115 in human, 133 in the mouse) -long TH2 cytokine that is part of the hematopoietic family. Unlike other members of this cytokine family (namely IL-3 and GM-CSF), this glycoprotein in its active form is a homodimer.

AA Sequence

MEIPMSTVVKETLIQLSTHRALLTSNETMRLPVPTHKN
HQLCIGEIFQGLDILKNQTVRGGTVEILFQNLSLIKKYI
DGQKEKCGEERRKTRHFLDYLQEFLGVMSTEWAME
V

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/ml. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Product Specification

Synonyms	B-cell Differentiation Factor I, Eosinophil Differentiation Factor, TRF
Accession	Q08125
Unigene	Rn.44227.
Species	Rat
Source	Escherichia coli.
Molecular Weight	Approximately 26.2 kDa, a disulfide-linked homodimeric protein containing two 113 amino acids.
Purity	> 98 % by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active when compared to standard. The ED ₅₀ as determined by a cell proliferation assay using human TF-1 cells is less than 0.5 ng/ml,

	corresponding to a specific activity of $> 2.0 \times 10^6$ IU/mg.
Physical Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation	Lyophilized from a 0.2 μ m filtered concentrated solution in PBS, pH 7.4.
Endotoxin	Less than 1 EU/ μ g of rRtIL-5 as determined by LAL method.