

Recombinant Rat Insulin-like Growth Factor-1

#R00325

Size: 10 µg/100 µg/500 µg

Stability & Storage

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.- 1 month, 2 to 8 °C under sterile conditions after reconstitution.

- 3 months, -20 to -70 °C under sterile conditions after reconstitution.



Warning: This product is intended for scientific research only. It is not to be used for medical or diagnostic procedures in humans or animals, nor as a food, cosmetic, or household product.

Background

The insulin-like growth factors (IGFs) belonged to the insulin gene family, are mitogenic polypeptide growth factors that stimulate the proliferation and survival of various cell types including muscle, bone, and cartilage tissue in vitro. The IGFs are similar by structure and function to insulin, but have a much higher growth-promoting activity than insulin. IGF-1 is produced primarily by the liver as an endocrine hormone as well as in target tissues in a paracrine/autocrine fashion. The production of IGF-1 is stimulated by growth hormone (GH) and can be retarded by undernutrition, growth hormone insensitivity, lack of growth hormone receptors, or failures of the downstream signaling pathway post GH receptor including SHP2 and STAT5B. Recombinant human IGF-1 are globular proteins containing 70 amino acids and 3 intra-molecular disulfide bonds. Mature rat IGF-1 shares 96 % a.a. sequence identity with human IGF-1

and exhibits cross-species activity.

AA Sequence

GPETLCGAELVDALQFVCGPRGFYFNKPTGYGSSIR
RAPQTGIVDECCFRSCDLRRLEMYCAPLKPTKSA

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Product Specification

Synonyms	Somatomedin C, IGF-I, IGF-IA, Mechano growth factor, MGF
Accession	P08025
GeneID	24482
Species	Rat
Source	Escherichia coli.
Molecular Weight	Approximately 7.7 kDa, a single non-glycosylated polypeptide chain containing 70 amino acid residues.
Purity	> 97 % by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active when compared to standard. The ED ₅₀ as determined by a cell proliferation assay using serum free human MCF-7 cells is less than 2 ng/ml, corresponding to a

	specific activity of $> 5.0 \times 10^5$ IU/mg.
Physical	Sterile Filtered White lyophilized
Appearance	(freeze-dried) powder.
Formulation	Lyophilized from a 0.2 μ m filtered concentrated solution in PBS, pH 7.4.
Endotoxin	Less than 1 EU/ μ g of rRtIGF-1 as determined by LAL method.