

Recombinant Human Growth Regulated Protein-gamma/CXCL3

#R00356

Size: 10 µg/100 µg/500 µg

Stability & Storage

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.



Warning: This product is intended for scientific research only. It is not to be used for medical or diagnostic procedures in humans or animals, nor as a food, cosmetic, or household product.

Background

CXCL3, also named GRO-γ, is belonging to the CXC chemokine family and encoded by the gene CXCL3. CXCL3/GRO-γ shares 86 % amino acid sequence with CXCL1/GRO-α. All three human GROs (GRO-α, GRO-β, GRO-γ) are members of the intercrine alpha (chemokine C-X-C) subfamily of chemokines. This chemokine is secreted by monocytes and macrophages. The functional receptor for CXCL3 has been identified as CXCR2. Similar to other GRO proteins, CXCL3 is potent neutrophil attractants and activators. CXCL3 plays a role in inflammation and exert its effects on endothelial cells in an autocrine fashion. All three GROs can bind with high affinity to the IL-8 receptor type B.

AA Sequence

ASVVTELRCQCLQTLQGIHLKNIQSVNVRSPGPHCA
QTEVIATLKNKGKKACLNPASPMVQKIIIEKILNKGSTN

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Product Specification

Synonyms	GRO-gamma, MIP2-beta
Accession	P19876
GeneID	2921
Species	Human
Source	Escherichia coli.
Molecular Weight	Approximately 7.9 kDa, a single non-glycosylated polypeptide chain containing 73 amino acids.
Purity	> 97 % by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human CXCR2 transfected human 293 cells is in a concentration range of 10-100 ng/ml.

Physical	Sterile Filtered White lyophilized
Appearance	(freeze-dried) powder.
Formulation	Lyophilized from a 0.2 μm filtered concentrated solution in 20 mM PB, pH 7.4, 50 mM NaCl.
Endotoxin	Less than 1 EU/μg of rHuGRO-γ/CXCL3 as determined by LAL method.