

Recombinant Human I-309/CCL1

#R00374

Size: 10 µg/100 µg/500 µg

Stability & Storage

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.



Warning: This product is intended for scientific research only. It is not to be used for medical or diagnostic procedures in humans or animals, nor as a food, cosmetic, or household product.

Background

Chemokine (C-C motif) ligand 1 (CCL1) belongs to a family inflammatory cytokines and also known as chemokines. It is a small glycoprotein secreted by activated T cells with a molecular weight of approximately 8.5 kDa. CCL1 attracts monocytes, NK cells, and immature B cells and dendritic cells by interacting with a cell surface chemokine receptor called CCR8. Human CCL1 has been assumed to be a homologue of the mouse TCA3. While the two proteins share only approximately 42 % amino acid sequence identity, both chemokines contain an extra pair of cysteine residues not found in most other chemokines.

AA Sequence

SKSMQVPFSRCCFSFAEQEIPLRAILCYRNTSSICSNE
GLIFKLKRGKEACALDTVGWVQRHRKMLRHCP SKRK

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Product Specification

Synonyms	Small-inducible cytokine A1, T lymphocyte-secreted protein I-309
Accession	P22362
GeneID	6346
Species	Human
Source	Escherichia coli.
Molecular Weight	Approximately 8.6 kDa, a single, non-glycosylated polypeptide chain containing 74 amino acids.
Purity	>97% by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human T-lymphocytes is in a concentration range of 10-100

	ng/ml.
Physical	Sterile Filtered White lyophilized
Appearance	(freeze-dried) powder.
Formulation	Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, pH 7.4, 100 mM NaCl.
Endotoxin	Less than 1 EU/µg of rHul-309/CCL1 as determined by LAL method.
