

Recombinant Human Monocyte Chemotactic Protein-1/CCL2

#R00375

Size: 5 µg/100 µg/500 µg

Stability & Storage

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.



Warning: This product is intended for scientific research only. It is not to be used for medical or diagnostic procedures in humans or animals, nor as a food, cosmetic, or household product.

Background

CCL2, also named monocyte chemotactic protein-1 (MCP-1) and small inducible cytokine A2, is a small cytokine that belongs to the CC chemokine family. It is a monomeric polypeptide, with a molecular weight of approximately 13 kDa and it is anchored in the plasma membrane of endothelial cells by glycosaminoglycan side chains of proteoglycans. Research shows that CCR2 and CCR4 are two cell surface receptors that bind CCL2. CCL2 exhibits a chemotactic activity for monocytes and basophils. However, it does not attract neutrophils or eosinophils. CCL2 has been implicated in the pathogenesis of diseases characterized by monocytic infiltrates, like psoriasis, rheumatoid arthritis or atherosclerosis and it may be involved in the recruitment of monocytes into the arterial wall during the disease process of atherosclerosis. Mature human CCL2 shares 78-79 % a.a. identity with canine, porcine and equine CCL2, while mouse and rat express a form

of CCL2 that is extended by 49 a.a. and shares only ~56 % a.a. identity within the common region.

AA Sequence

QPDAINAPVTCCYNFTNRKISVQRLASYRRITSSKCP
KEAVIFKTIVAKEICADPKQKWVQDSMDHLDKQTQTP
KT

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Product Specification

Synonyms	HC11, Monocyte chemoattractant protein 1, MCAF, MCP-1, Monocyte secretory protein JE, Small-inducible cytokine A2
Accession	P13500
GeneID	6347
Species	Human
Source	Escherichia coli.
Molecular Weight	Approximately 8.7 kDa, a single non-glycosylated polypeptide chain containing 76 amino acids.
Purity	> 96 % by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using

human monocytes is in a concentration range of 10-100 ng/ml.

Physical Appearance Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, pH 7.4, 100 mM NaCl.

Endotoxin Less than 1 EU/µg of rHuMCP-1/CCL2 as determined by LAL method.
