

Recombinant Human Thymus and Activation Regulated Chemokine/CCL17

#R00390

Size: 5 µg/100 µg/500 µg

Stability & Storage

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.



Warning: This product is intended for scientific research only. It is not to be used for medical or diagnostic procedures in humans or animals, nor as a food, cosmetic, or household product.

Background

Human CCL17 also known as thymus and activation-related chemokine (TARC) is encoded by the CCL17 gene located on the chromosome 16 in humans. It is expressed by thymus cells constitutively and phytohemagglutinin-stimulated peripheral blood mononuclear cells transiently. CCL17 signals through the chemokine receptors CCR4 and CCR8 and displays chemotactic activity for T lymphocytes and some other leukocytes. It plays an important role in skin diseases such as atopic dermatitis, bullous pemphigoid and mycosis fungoides. CCL17 has approximately 24 – 29 % amino acid sequence identity with RANTES, MIP-1α, MIP-1β, MCP-1, MCP-2, MCP-3 and I-309.

AA Sequence

ARGTNVGRECCLEYFKGAIPRLRKLKTWYQTSEDCSR
DAIVFVTVQGRAICSDPNNKRVKNAVKYLQSLERS

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Product Specification

Accession	Q92583
GeneID	6361
Species	Human
Source	Escherichia coli.
Molecular Weight	Approximately 8.1 kDa, a single non-glycosylated polypeptide chain containing 71 amino acids.
Purity	> 97 % by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human T-lymphocytes is in a concentration range of 1.0-10 ng/ml.
Physical Appearance	Sterile Filtered White Lyophilized (freeze-dried) powder.

Formulation	Lyophilized from a 0.2 μm filtered concentrated solution in 20mM PB, pH 7.4, 150mM NaCl.
Endotoxin	Less than 1 EU/μg of rHuTARC/CCL17 as determined by LAL method.