

Recombinant Murine Macrophage Inflammatory Protein-2/CXCL2

#R00409

Size: 5 µg/100 µg/500 µg

Stability & Storage

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.



Warning: This product is intended for scientific research only. It is not to be used for medical or diagnostic procedures in humans or animals, nor as a food, cosmetic, or household product.

Background

Murine CXCL2, also named MIP-2, is belonging to the CXC chemokine family. It is encoded by the gene CXCL2. CXCL2 shares 90 % amino acid sequence with CXCL1/GROα. It is member of the intercrine alpha (chemokine C-X-C) subfamily of chemokines. This chemokine is secreted from a murine macrophage cell line. The functional receptor for CXCL2 has been identified as CXCR2. CXCL2 is chemotactic for polymorphonuclear leukocytes and hematopoietic stem cells. Similar to other GRO proteins, CXCL2 is potent neutrophil attractants and activators. In addition, it is also active toward basophils. The amino acid sequence of murine CXCL2 is 60 % identical to human CXCL2.

AA Sequence

AVVASELRCQCLKTLPRVDFKNIQSLSVTPPGPHCAQ
TEVIATLKGGQKVCLEAPLVQKIIQKILNKGKAN

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Product Specification

Accession	P10889
GeneID	20310
Species	Murine
Source	Escherichia coli.
Molecular Weight	Approximately 7.8 kDa, a single, non-glycosylated polypeptide chain containing 73 amino acids.
Purity	> 97 % by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human neutrophils is in a concentration range of 1.0-10 ng/ml.
Physical Appearance	Sterile Filtered White Lyophilized (freeze-dried) powder.

Formulation	Lyophilized from a 0.2 μm filtered concentrated solution in 20 mM PB, pH 7.4, 150 mM NaCl.
Endotoxin	Less than 1 EU/μg of rMuMIP-2/CXCL2 as determined by LAL method.