

Recombinant Murine I-TAC/CXCL11

#R00416

Size: 5 µg/100 µg/500 µg

Stability & Storage

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.



Warning: This product is intended for scientific research only. It is not to be used for medical or diagnostic procedures in humans or animals, nor as a food, cosmetic, or household product.

AA Sequence

FLMFKQGRCLCIGPGMKAVKMAEIEKASVIYPSNGCD
KVEVIVTMKAHKRQRCLDPRSKQARLIMQAIEKKNFL
RRQNM

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Product Specification

Accession	Q9JHH5
GeneID	56066
Species	Murine
Source	Escherichia coli.
Molecular Weight	Approximately 9.1 kDa, a single non-glycosylated polypeptide chain containing 79 amino acids.
Purity	> 98 % by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using murine CXCR3 transfected 293 cells is in a concentration of 10-100 ng/ml.
Physical Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation	Lyophilized from a 0.2 µm filtered concentrated solution in 10 mM Sodium Citrate, pH 4.0, with 600 mM NaCl.
Endotoxin	Less than 1 EU/µg of rMuI-TAC/CXCL11 as determined by LAL method.