

Recombinant Murine Stromal-Cell Derived Factor-1 alpha/CXCL12α

#R00417

Size: 10 µg/100 µg/500 µg

Stability & Storage

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.



Warning: This product is intended for scientific research only. It is not to be used for medical or diagnostic procedures in humans or animals, nor as a food, cosmetic, or household product.

Background

CXCL12 also known as SDF-1 is belonging to the CXC chemokine family. It is encoded by the CXCL12 gene. Murine CXCL12 is expressed as two isoforms that differ only in the C-terminal tail. And both SDF-1 isoforms undergo proteolytic processing of the first two N-terminal amino acids. Contrast to SDF-1β, SDF-1α is shorter by four amino acids at the C-terminal tail. On the cell surface, the receptor for this chemokine is CXCR4 and syndecan4. CXCL12 is strongly chemotactic for T-lymphocytes, monocytes, but not neutrophils. SDF-1 is highly conserved between species, murine CXCL12α shares approximately 93% amino acid sequence identity with human CXCL12α.

AA Sequence

KPVSLSYRCPCRFFESHIANVKHLKILNTPNCALQI
VARLKNNNRQVCIDPKLKWIQEYLEKALNK

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Product Specification

Synonyms	TPAR1, PBSF, TLSF
Accession	P40224
GeneID	20315
Species	Murine
Source	Escherichia coli.
Molecular Weight	Approximately 8.0 kDa, a single non-glycosylated polypeptide chain containing 68 amino acids.
Purity	> 97 % by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human peripheral blood monocytes is in a concentration range of 50-100 ng/ml.
Physical	Sterile Filtered White Lyophilized

Appearance	(freeze-dried) powder.
Formulation	Lyophilized from a 0.2 μm filtered concentrated solution in PBS, pH 7.4.
Endotoxin	Less than 1 EU/μg of rMuSDF-1α/CXCL12α as determined by LAL method.