

Recombinant Murine Fractalkine/CX3CL1

#R00423

Size: 5 µg/100 µg/500 µg

Stability & Storage

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.CHAR(10)- 12 months from date of receipt, -20 to -70 °C as supplied.CHAR(10)- 1 month, 2 to 8 °C under sterile conditions after reconstitution.CHAR(10)- 3 months, -20 to -70 °C under sterile conditions after reconstitution.



Warning: This product is intended for scientific research only. It is not to be used for medical or diagnostic procedures in humans or animals, nor as a food, cosmetic, or household product.

AA Sequence

QHLGMTKCEIMCGKMTSRIPVALLIRYQLNQESCGKR
AIVLETTQHRRFCADPKEKWVQDAMKHLDHQAAALT
KNG

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Product Specification

Accession	O35188
GeneID	20312
Species	Murine
Source	Escherichia coli.
Molecular Weight	Approximately 8.7 kDa, a single non-glycosylated polypeptide chain containing 76 amino acids and comprises only the chemokine domain of Murine Fractalkine.
Purity	> 97 % by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active when compared to standard. The ED ₅₀ as determined by a cell proliferation assay using human peripheral blood lymphocytes (PBL) is less than 0.5 µg/ml,

	corresponding to a specific activity of > 2000 IU/mg.
Physical Appearance	Sterile Filtered White Lyophilized (freeze-dried) powder.
Formulation	Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.
Endotoxin	Less than 0.1 EU/µg of rMuFractalkine/CX3CL1 as determined by LAL method.