

Recombinant Murine Macrophage Inflammatory Protein-1 gamma/CCL9/CCL10

#R00430

Size: 5 µg/100 µg/500 µg

Stability & Storage

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.



Warning: This product is intended for scientific research only. It is not to be used for medical or diagnostic procedures in humans or animals, nor as a food, cosmetic, or household product.

Background

Murine CCL9/10 is belonging to the CC chemokine family. It is encoded by the gene CCL9. MRP-2 (CCL9/10) was isolated from a murine macrophage cell line, RAW 264.7. MRP-2 shows a 51 % sequence identity at the protein level to MRP-1 (C10) and 46% identity to MIP-1-alpha (CCL3). The functional receptor for CCL9/10 has been identified as CCR1. In contrast, other CCR1 ligands, CCL3/MIP-1alpha and CCL5/RANTES are considered to be primary CCR1 ligands in inflammatory responses. CCL9 can activate osteoclasts suggesting an important role for it in bone resorption. Murine CCL9/10 is 75 % amino acid identical to rat CCL9/10.

AA Sequence

QITHATETKEVQSSLKAQQGLEIEMFHMGMFQDSSDC
CLSYNSRIQCSRFIGYFPTSGGCTRPGIIFISKRGFQV
CANPSDRRVQRCIERLEQNSQPRTYKQ

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Product Specification

Synonyms	MIP-1-gamma, MRP-2, Small-inducible Cytokine A9
Accession	P51670
GeneID	20308
Species	Murine
Source	Escherichia coli.
Molecular Weight	Approximately 11.6 kDa, a single non-glycosylated polypeptide chain containing 101 amino acids.
Purity	> 95 % by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human neutrophils is in a concentration range of 0.1-10

	ng/ml.
Physical	Sterile Filtered White lyophilized
Appearance	(freeze-dried) powder.
Formulation	Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.
Endotoxin	Less than 1 EU/µg of rMuMIP-1γ/CCL9/CCL10 as determined by LAL method.