

Recombinant Rat Neutrophil Activating Peptide 2/CXCL7

#R00448

Size: 10 µg/100 µg/500 µg

Stability & Storage

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.



Warning: This product is intended for scientific research only. It is not to be used for medical or diagnostic procedures in humans or animals, nor as a food, cosmetic, or household product.

Background

CXCL7 is a small cytokine which is also an isoform of Beta-Thromboglobulin or Pro-Platelet basic protein belonging to the CXC chemokine family and it is released in large amounts from platelets following their activation. CXCL7 stimulates DNA synthesis, mitosis, glycolysis, intracellular cAMP accumulation, prostaglandin E2 secretion, and synthesis of hyaluronic acid and sulfated glycosaminoglycan. It also stimulates the formation and secretion of plasminogen activator by human synovial cells. Recombinant Rat CXCL7 contains 62 amino acids which is a single non-glycosylated polypeptide chain. In addition, The rat CXCL7 shares 58 % and 77 % a.a. sequence identity with human and mouse CXCL7.

AA Sequence

IELRCRCTNTLSGIPLNSISRVNVFRPGAHCNDNVEVIA
TLKNGKEVCLDPTAPMIKKIVKKI

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/ml. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Product Specification

Synonyms	CXC Chemokine RTCK1, Chemokine (C-X-C Motif) Ligand 7, Isoform CRA_b, Protein Ppbp.
Species	Rat
Source	Escherichia coli.
Molecular Weight	Approximately 6.8 kDa, a single non-glycosylated polypeptide chain containing 62 amino acids.
Purity	> 97 % by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active when compared to standard. The biologically active determined by a chemotaxis bioassay using human CXCR2 transfected murine BaF3 cells is in a concentration range of 0.1-1.0 ng/ml.

Physical	Sterile Filtered White lyophilized
Appearance	(freeze-dried) powder.
Formulation	Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.
Endotoxin	Less than 1 EU/µg of rRtNAP-2/CXCL7 as determined by LAL method.