

Recombinant Rat gamma-Interferon Inducible Protein 10/CXCL10

#R00449

Size: 5 µg/100 µg/500 µg

Stability & Storage

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.



Warning: This product is intended for scientific research only. It is not to be used for medical or diagnostic procedures in humans or animals, nor as a food, cosmetic, or household product.

Background

CXCL10 also known as IP-10 is belonging to the CXC chemokine family. It is encoded by the CXCL10 gene, and in rat it is also named the Mob-1 gene. This gene was identified on the basis of differential expression in rat embryo fibroblasts. CXCL10 elicits its effects by binding to the cell surface chemokine receptor CXCR3. CXCL10 has been shown to be a chemoattractant for activated T-lymphocytes and monocytes/macrophages. It also has other roles, such as promotion of T cell adhesion to endothelial cells, and inhibition of bone marrow colony formation and angiogenesis. Rat CXCL10 shares approximately 68 % amino acid sequence identity with human CXCL10.

AA Sequence

IPLARTVRCTCIDFHEQPLRPRAIGKLEIIPASLSCPHV
EIIATMKNNNEKRCLNPESEAIKSLKAVSQRRSKRAP

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Product Specification

Synonyms	Gamma-IP10, Small-inducible Cytokine B10, Protein Mob-1
Accession	P48973
GeneID	245920
Species	Rat
Source	Escherichia coli.
Molecular Weight	Approximately 8.7 kDa, a single non-glycosylated polypeptide chain containing 77 amino acids.
Purity	> 95 % by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human CXCR3 transfected HEK293 cells is in a concentration range of 10-50

	ng/ml.
Physical	Sterile Filtered White lyophilized
Appearance	(freeze-dried) powder.
Formulation	Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.
Endotoxin	Less than 1 EU/µg of rRtIP-10/CXCL10 as determined by LAL method.