

Recombinant Human Melanoma Inhibitor Activity Protein 2

#R00539

Size: 5 µg/100 µg/500 µg

Stability & Storage

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.



Warning: This product is intended for scientific research only. It is not to be used for medical or diagnostic procedures in humans or animals, nor as a food, cosmetic, or household product.

Background

Melanoma Inhibitor Activity Protein 2 (MIA-2) encoded by MIA2 gene in humans, is a secreted cytokine that is highly expressed in liver and weakly in testis. The patients with severe fibrosis or inflammation and chronic hepatitis have higher level of MIA2 than other. Levels of MIA2 may severe as a clinically utility marker for diagnosis of hepatic disease activity and severity. The MIA2 is a member of the MIA/OTOR family, which also includes MIA, OTOR, and TANGO, and they share a Src homology-3 (SH3)-like domain.

AA Sequence

LESTKLLADLKKCGDLECEALINRVSAMRDYRGPD
CR YLNFTKGEEISVYVKLAGEREDLWAGSKGKEFGYFP
RDAVQIEEVFISEEIQMSTKESDFLCL

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom.

Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Product Specification

Accession	Q96PC5
GeneID	117153
Species	Human
Source	Escherichia coli.
Molecular Weight	Approximately 11.4 kDa, a single non-glycosylated polypeptide chain containing 100 amino acids.
Purity	> 97 % by SDS-PAGE and HPLC analyses.
Biological Activity	Data Not Available.
Physical Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation	Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, pH 7.4, 150 mM NaCl.
Endotoxin	Less than 1 EU/µg of rHuMIA2 as determined by LAL method.